


Very Long (Effective Area: 127 × 4mm)

This new long-scale MCP allows for continuous detection of wide range position information without dead space correction. This new configuration promises reliable measurement, and is well suited for double-focusing mass spectrometers. Assembled configurations are also available upon request.

FEATURES

- Long Scale (Effective Area: 127mm × 4mm)
- Simultaneous Multielement Measurement Possible
- Sensitive to Ions, Electrons, UV Radiation, Soft X-rays, γ -rays and High Energy Particles
- High Gain (1×10^4)



TMCPF0064

APPLICATIONS

- SIMS (Secondary Ions Mass Spectrometers)
- UV Spectrometers
- X-ray Spectrometers
- Electron Spectrometers

GENERAL

Parameter	Description/Value	Unit
Outer Dimension	140 × 9	mm
Electrode Dimension	138 × 8	mm
Effective Area Dimension	127 × 4	mm
Thickness	0.48	mm
Channel Diameter	12	μm
Channel Pitch	15	μm
Bias Angle	8	$^\circ$
Open Area Ratio	60	%
Electrode Material	Inconel	—

CHARACTERISTICS (at 1000V, 1.3×10^{-4} Pa (1×10^{-6} Torr), 25°C)

Parameter	Description/Value	Unit
Current Gain Min.	1×10^4	—
Plate Resistance	10 to 100	$\text{M}\Omega$
Maximum Dark Current	5×10^{-13}	A/cm^2
Maximum Linear Output Current	up to 7% of the strip current [Ⓐ]	—


MAXIMUM RATINGS (Absolute Values)

Parameter	Description/Value	Unit
Supply Voltage [Ⓑ]	1000	V
Ambient Temperature	-50 to +50	$^\circ\text{C}$

NOTE: Ⓐ: Strip current is current flowing through the channel walls, which supplies the current released from the channel walls. It is given by: Supply voltage / Plate resistance.


Ⓑ: At a vacuum of 1.3×10^{-4} Pa (1×10^{-6} Torr) or better.

Figure 1: Typical Current Gain




LONG SCALE MICROCHANNEL PLATE F6492

Figure 2: Dimensional Outline (Unit: mm)



TMCPA0023EB

Figure 3: Application Example to SIMS¹⁾



TMPC0038EA

PRECAUTIONS FOR USE

- Avoid touching the MCP or MCP assembly with bare hands.
- Handle the MCP only in a clean room since dust and humidity may adversely affect MCP characteristics.
- The MCP should be operated in vacuum below 1.33×10^{-4} Pa (1×10^{-6} torr).
- The MCP should be kept in vacuum or dry nitrogen gas atmosphere during long periods of storage.
- When outgassing from the MCP occurs, baking the MCP at 350°C maximum in a vacuum system is recommended. In addition electron bombarding may be effective.

REFERENCE

- 1) Japan Academic Promotion Society, 141th Committee on Micro Beam Analysis: "Micro beam analysis" Asakura Shoten, 293 (1985)

HAMAMATSU

HAMAMATSU PHOTONICS K.K., Electron Tube Center

314-5, Shimokanzo, Toyooka-village, Iwata-gun, Shizuoka-ken, 438-0193, Japan, Telephone: (81)539/62-5248, Fax: (81)539/62-2205

U.S.A.: Hamamatsu Corporation: 360 Foothill Road, Bridgewater. N.J. 08807-0910, U.S.A., Telephone: (1)908-231-0960, Fax: (1)908-231-1218

Germany: Hamamatsu Photonics Deutschland GmbH: Arzbergerstr. 10, D-82211 Herrsching am Ammersee, Germany, Telephone: (49)8152-375-0, Fax: (49)8152-2658

France: Hamamatsu Photonics France S.A.R.L.: 8, Rue du Saule Trapu, Parc du Moulin de Massy, 91882 Massy Cedex, France, Telephone: (33)1 69 53 71 00, Fax: (33)1 69 53 71 00

United Kingdom: Hamamatsu Photonics UK Limited: Lough Point, 2 Gladbeck Way, Windmill Hill, Enfield, Middlesex EN2 7JA, United Kingdom, Telephone: (44)181-367-3560, Fax: (44)181-367-6384

North Europe: Hamamatsu Photonics Norden AB: Färögatan 7, S-164-40 Kista Sweden, Telephone: (46)8-703-29-50, Fax: (46)8-750-58-95

Italy: Hamamatsu Photonics Italia: S.R.L.: Via Della Moia, 1/E, 20020 Arese, (Milano), Italy, Telephone: (39)2-935 81 733, Fax: (39)2-935 81 741

TMCP1009E05
JAN. 1997 SI (9609)
Printed in Japan